



Mobilizing the immune system
to fight inflammatory and viral
diseases, as well as cancer

Abivax, a late-stage clinical biotech company

July, 2020



Forward looking statements

This presentation contains information pertaining to Abivax S.A. (“Abivax”). Neither Abivax, nor its management, shareholders, directors, advisors, employees or representatives make any representation or warranty, express or implied, as to the fairness, the accuracy, completeness or correctness of any information contained in this presentation or any other information transmitted or made available to the viewer or recipient hereof, whether communicated in written or oral form. Neither Abivax, nor its management, shareholders, directors, advisors, employees or representatives accept any responsibility in this respect.

This presentation contains forward-looking statements. These statements reflect management’s current views with respect to Abivax’s product candidates’ development, clinical and regulatory timelines and anticipated results, market opportunity, potential financial performance and other statements of future events or conditions, which are naturally subject to risks and contingencies that may lead to actual results materially differing from those explicitly or implicitly included in these statements. Although Abivax believes that the expectations reflected in such forward-looking statements are reasonable, no assurance can be given that such expectations will prove to have been correct. Accordingly, results could differ materially from those set out in the forward-looking statements as a result of various factors, many of which are beyond Abivax’s control. No reliance should be made on such forward-looking statements.

Abivax does not undertake to update or revise the presentation, including the forward-looking statements that may be presented in this document to reflect new information, future events or for any other reason, following distribution, beyond what is required by applicable law or applicable stock exchange regulations if and when circumstances arise that will lead to changes compared to the date when these statements were provided.

In the European Union (including in France), this presentation is intended solely for “qualified investors” within the meaning of Article 2(1)(e) of the Prospectus Directive (Directive 2003/71/EC) as amended (including amendments by Directive 2010/73/EU), to the extent implemented in the relevant member state). This presentation has been prepared on the basis that any offering of securities by the Company in any member state of the European Economic Area has implemented the Prospectus Directive (2003/71/EC) will be made either by means of a prospectus filed with the authority of the relevant member state, or pursuant to an exemption under the Prospectus Directive, as implemented in that relevant member state, from the requirement to publish a prospectus.

This presentation does not constitute or form part of, and should not be construed as, an offer to sell or issue or the solicitation of an offer to buy or acquire securities of Abivax, in any jurisdiction or an inducement to enter into investment activity, nor shall there be any sale of securities in any state or jurisdiction in which such offer, solicitation or sale would be unlawful prior to registration or qualification under the securities law of any such state or jurisdiction. No part of this presentation, nor the fact of its distribution, should form the basis of, or be relied on in connection with any contract or commitment or investment decision whatsoever.

Key company facts

Milestones



Founded in 2013
by Truffle Capital



Sept. 2018: Focus ABX464
on chronic inflammation



Abivax went
public in June 2015,
raising € 57.7m



May 2020: ABX464 to
treat acute viral and
inflammatory diseases

Location



Head Office
Paris


**Cooperative
Lab with CNRS**
Montpellier



BREAKING NEWS

**Abivax treats first patient
in Phase 2b/3 ABX464
COVID-19 clinical trial**

Shareholder structure¹ and market cap²



■ Truffle Capital
■ Sofinnova
■ Board & management
■ Incubator & founders
■ Public

Operations



26
Employees



Cash³
€ 9.8m




20
in R&D




6
in Support

- 1) Undiluted – as of 31.03.2020
2) As of 22.05.2020 EOB
3) Actual December 2019

Abivax: A strong and diversified pipeline



ABX464: A promising candidate addressing attractive markets



Total market size*
in inflammatory
diseases

greater than
USD 90 B



Market size*
in first indication
(ulcerative colitis)

around
USD 5.8 B

Coming from the **proprietary** Abivax library of compounds, biased to **modulate RNA biogenesis** (>2200 molecules); Close collaboration with EVOTEC

Small molecule (quinoline), administered as an **oral capsule** (once a day)

First-in-Class, novel mechanism of action: Selective upregulation of anti-inflammatory microRNA miR-124

Good safety profile after administration to **>300 patients and volunteers**


Anti-inflammatory effect confirmed in DSS mouse model of IBD as well as in Phase 2a induction and maintenance studies in ulcerative colitis. Phase 2b study in UC ongoing in 232 Patients, as well as Phase 2a in rheumatoid arthritis in 60 patients

High medical need for novel safe and efficacious drugs in inflammatory diseases

* For Europe G5, U.S. and Japan

ABX464-101/102 study design: Phase 2a in ulcerative colitis

- Randomized, double-blind, placebo controlled, multi-national study followed by an open-label maintenance study



Study Population

- Moderate to severe active UC patients who **failed or were intolerant to immunomodulators, anti-TNF α , vedolizumab and/or corticosteroids**
- **Confirmed UC** for at least 3 months with a **Total Mayo Score of 6–12** with **endoscopic sub-score of 2 or 3**

Central reading of endoscopies (Induction)

ABX464-101:

Statistically significant efficacy achieved for major endpoints (day 56)

Clinical remission:

Total Mayo Score (TMS) equal or lower than 2 + no sub-score >1

Endoscopic improvement:

Endoscopy sub-score 0 or 1

Clinical response:

TMS decrease of min 3 points and 30% from baseline + decrease of bleeding sub-score of min 1 point or absolute baseline of 0 or 1



	ABX464 (n=20/23) PP/ITT	Placebo (n=9/9) PP/ITT	p value (PP)
Clinical remission*	35%/30%	11%/11%	0.16
Endoscopic improvement	50%/43%	11%/11%	0.03
Clinical response	70%/61%	33%/33%	0.06
Total Mayo Score reduction	-53%	-27%	0.03
Partial Mayo Score reduction	-62%	-32%	0.02
miR-124 expression in rectal biopsies (fold increase)	7.69	1.46	0.004


*Clinical remission according to previous FDA definition. With application of most recent FDA definition (excluding physician assessment), clinical remission rate was 40% in ABX464 group and remained at 11% with placebo

ABX464-101 Partial Mayo Score Results

Fast onset of action and clear responses in patients previously treated with biologics


Overall Patient Population

Change from Baseline Partial Mayo Score



Patients previously treated with biologics

Change from Baseline Partial Mayo Score



ABX464-102:

Efficacy further amplified by 12-months maintenance study

29/32

Patients completed the induction study

4/6

Countries granted regulatory approval for maintenance study

22/23

Eligible patients enrolled in the maintenance study, 19 completed first year


Fast onset and durable efficacy with impressive 12 months data.

	Day 0 Maintenance	Month 12
Clinical remission (TMS) (no loss of clinical remission)	6/19 (31.6%)	12/16 (75.0%)*
Clinical Response	14/19 (73.7%)	15/16 (93.7%)
Endoscopic Improvement** (sub-score 0 or 1)	11/19 (57.9%)	16/16 (100.0%)
Endoscopic remission*** (sub-score 0)	2/19 (10.5%)	13/16 (81.2%)


As of June, 2020: Mean cumulative exposure to ABX464 in maintenance study is over two years with ABX464-102 UC maintenance patients N=16 (mean: 25.6 months; min.: 23.3 months; max: 30.5 months)

Changes of Partial Mayo Score and fecal calprotectin during the maintenance phase for all patients and patients previously on biologics

Partial Mayo Score – Mean



Fecal calprotectin $\mu\text{g/g}$ – Median



— All patients (N=19) — Patients previously treated with biologics (N=7)

→ Partial Mayo Score continued to decrease

→ Fecal calprotectin levels went down to normal values ($< 50 \mu\text{g/g}$)

ABX464, Vedolizumab, Tofacitinib and Filgotinib efficacy in induction and maintenance clinical trials

	Vedoluzimab Phase 3			Tofacitinib Phase 3			Filgotinib* Phase 3			ABX464 Phase 2a		
INDUCTION	Active	Placebo	Delta	Active	Placebo	Delta	Active	Placebo	Delta	Active	Placebo	Delta
Clinical Remission (%)	16,9	5,4	11,5	16,8-18,5	3,6-8,2	13,2-10,3	11,5-26,1**	4,2-15,3	7,3-10,8	35	11	24
Mucosal Healing (%)	40,9	24,8	16,1	28,4-31,3	11,6-15,6	16,8-15,7	n/a	n/a	n/a	50	11	39
MAINTENANCE												
Clinical Remission (%)	41,8	15,9	25,9	34,3-40,6	11,1	23,2-29,5	37,2	11,2	26	75		
Mucosal Healing (%)	51,6	19,8	31,8	37,4-45,7	13,1	24,3-32,6	n/a	n/a	n/a	100		

* For patients treated with 200mg

** Biologic experienced vs. biologic naïve patients

Phase 2a study ABX464-102 maintenance study allowed all patients irrespective of treatment assignment or clinical response during induction to be included in open label ABX464 50mg open label

ABX464 showed a good safety profile during clinical development of the 50 mg dosage form

Safety profile consistent with previous and ongoing clinical studies

(>300 healthy volunteers and patients exposed to ABX464)



Overall: Generally well tolerated with no deaths, no malignancies, no severe infections, no significant changes in the laboratory parameters including blood cell counts

No Serious Adverse Reactions, most AEs were of mild to moderate intensity

Most frequently reported AEs: Headache and epigastric pain; occurring mainly during the first days of treatment

N=16 patients from UC ABX464-102 maintenance study are on **continuous daily treatment with ABX464 with a mean of 26 months**

Conclusions

ABX464 oral 50mg QD drug candidate for moderate to severe UC patients



Good safety and tolerability of chronic treatment with ABX464 50mg QD in patients with UC

Conclusion is supported by safety analysis in more than 300 healthy volunteers and patients (no serious adverse reactions, no severe infections, no lymphopenia, no neutropenia)

Most frequently reported adverse events were transient and mild headache, nausea, gastro-intestinal pain



Confirmed efficacy in Phase 2a UC induction study

- All endpoints favorable to ABX464, with statistical significance in endoscopic improvement, TMS and PMS reductions, and clear trends for clinical remission and clinical response
- Fast onset of action
- Active in both biologics naive and biologics refractory patients



Efficacy signal further amplified during 12-months maintenance study

- Continued very good safety profile
- Durability of clinical efficacy with further improvement and increased clinical remission with longer treatment
- Normalized fecal calprotectin levels
- Significant endoscopic improvement
- Continued over expression of miR-124

ABX464 ongoing and planned studies



Phase 2b in ulcerative colitis:

- Conducted with IQVIA as CRO
- 232 patients, 17 countries, 150+ study sites
- 4 study arms (placebo, 25, 50, 100 mg QD)
- Central blinded reading of endoscopies
- Top-line data for induction phase expected for Q2 2021



Phase 2b study in 232 patients with moderate to severe ulcerative colitis is currently ongoing in Canada and Europe and the US (with FDA clearance of IND on 19/01/20)



Phase 2a study ongoing in 60 patients with rheumatoid arthritis in 5 European countries




Phase 2b study planned in app. 200 patients with Crohn's disease – FPI planned for end of 2020




Phase 2b/3 study in COVID-19 ongoing – FPI 01/07/2020, 2020, 1.034 patients in total

COVID-19 infection and pathology



ABX464 COVID-19 Development Rationale



Antiviral: ABX464 inhibits SARS-CoV-2 (COVID-19 virus) *in vitro* replication in human respiratory epithelium: Inhibition of COVID-19 viral replication comparable to Remdesivir

Anti-inflammatory: ABX464 has demonstrated potent anti-inflammatory properties in several *in-vivo* models and in patients with moderate to severe ulcerative colitis


Tissue repair observed in DSS model of inflammatory bowel disease (IBD) and in patients in Phase 2 ulcerative colitis trial

Good safety profile of ABX464 demonstrated in >300 patients and volunteers

Manufacturing capacity in place (drug substance, finished product and packaging) to supply the investigational drug for large clinical trials and rapidly scale for commercial production


ABX464 rapidly upregulates miR-124 (10-fold) within 24 hours in human PBMCs (*in vitro* results)

miR-124 upregulation vs. time of ABX464 treatment



ABX464 novel mechanism of action: Potent and specific upregulation of miR-124 leads to reduction of pro-inflammatory cytokines

Both systemic and local inflammatory sites



Established miR-124 targets:
(translation ↓)

Outcome

MCP-1/
CCL2

MCP-1/CCL2



STAT 3

IL-6

IL-17

TNF α



IL-6R

JAK pathway
blocked


IL-6

TNF- α



Anti-inflammatory effect: ABX464 showed efficacy in the DSS mouse model*


ABX464 protects mice from death in the DSS mouse model




— No treatment (n=8)

— ABX464 60 days (n=8)

DSS without ABX464 leads to intestinal damage



ABX464 protects intestinal structure




→ **ABX464 reduced the expression of pro-inflammatory cytokines in colon tissue:**
IL-6 (2x), TNF (7.5x) and MCP-1 (6x)

*Chebli et al, Nature Scientific Reports 7: 4860 (2017)


Tissue repair in an ABX464 treated UC patient

Courtesy of Prof. Severine Vermeire

Before ABX464




After ABX464



Antiviral effect: Reduction of COVID-19 replication in an *in vitro* reconstituted human airway epithelial model

Infectious titrations TCID50 at 48 hours post infection



Comparable efficacy between Remdesivir and ABX464

ABX464 showed a good safety profile during clinical development of the 50 mg dosage form

Safety profile consistent with previous and ongoing clinical studies

(>300 healthy volunteers and patients exposed to ABX464)



Overall: Generally well tolerated with no deaths, no malignancies, no severe infections, no significant changes in the laboratory parameters including blood cell counts

No Serious Adverse Reactions, most AEs were of mild to moderate intensity


Most frequently reported AEs: Headache and epigastric pain; occurring mainly during the first days of treatment

N=16 patients from UC ABX464-102 maintenance study are on **continuous daily treatment with ABX464 with a mean of 26 months**

Phase 2b/3 clinical trial miR-AGE: High-risk patients, PRIOR to respiratory distress

- **Early treatment** of high-risk patients infected with COVID-19
- **Main objective:** A Phase 2b/3, randomized, double blind, placebo-controlled study of ABX464 to treat inflammation and prevent acute respiratory failure
- **Inclusion criteria:** COVID-19 patients aged ≥ 65 and aged ≥ 18 with at least one additional risk factor who are infected with SARS-CoV-2
- **Target population:** hospitalized and non-hospitalized patients
- **Main evaluation criterion:** Absence of high-flow oxygen (>3 l/min), assisted ventilation (positive pressure or intubation) and/or death after 28 days
- **Treatment duration:** 28 days
- **1,034 patients** will be included in **50 clinical study sites in Europe and South America**
 - ❖ Placebo + SOC group: 344 patients
 - ❖ ABX464 + SOC group: 690 patients (2 to 1 randomization)
 - ❖ Expected response rates: 75% on placebo, 83 % on ABX464 (alpha 0.05, beta 80%)
- **Parexel** selected as CRO; total study costs € 16m


ABX464 supply available for COVID-19 clinical trials and scalable for commercialization



Abivax Communication Plan until mid-2021

	Q2 2020	Q3 2020	Q4 2020	Q1 2021	Q2 2021
UC Phase 2a (ABX464)			2-years maintenance data		
UC Phase 2b (ABX464)			Enrollment completed		Top line results
RA Phase 2a (ABX464)			Enrollment completed		Top line results
Crohn's Phase 2b (ABX464)			FPI		
COVID-19 Phase 2b/3 (ABX464)	FPI 	Enrollment completed	Top line results		MAA/NDA submission
HCC Phase 1/2 (ABX196)			Enrollment completed (Dose escalation)	Top line results Dose escalation phase	

Abivax financial planning – Bpifrance € 36m funding first building block



The amount of € 36m will be paid within the next 12 months

Total amount of € 36m is made of € 20,1m grant (non-refundable) and € 15,9m loan (refundable when ABX464 is reaching commercial stage)

Total amount of € 36m is funding miR-AGE study as well as additional costs for ABX464 development and manufacturing scale up, required for potential ABX464 MAA (marketing authorization application) in COVID-19 by mid-2021

Abivax Financing

07/2020

06/2021

ABX464
Manufacturing scale-up

ABX464
Preclinical

ABX464
Regulatory clinical
studies

Abivax
New research

Abivax
G&A

Funding Need
€ 80m

ABX464 in UC: Phase 2a M & Phase 2b I+M*

ABX464 in CD: Phase 2b I+M

ABX464 in RA: Phase 2a I+M & Phase 2b I

ABX464 in COVID-19: Phase 2b/3

ABX196 in HCC: Phase 1/2

COMPLETED

- Bpifrance funding € 36m
- Kepler Cheuvreux Equity line € 11m
- Société Générale PGE € 5m

Cash runway until early 2021

€ 52m

€ 28m
or more

TO BE COMPLETED by summer 2020

Core plan

- Non-dilut. Public funding
- Venture loan
- Capital raise

Cash runway > mid 2021

Upside plan

- Partnering
- ABX464 COVID-19 commercialization

Next funding milestone planned in Q2 2021: Abivax Partnering AND/OR Abivax Nasdaq listing

* I: Induction phase; M: Maintenance phase

Abivax Analyst Reports Overview

Analyst	Country	Last update	Target Price	Recommendation
Bryan, Garnier & Co	France	03/12/2019	€ 37.50	Buy
Degroof Petercam	Belgium	13/12/2019	€ 29.00	Buy
goetzpartners securities	UK/Germany	14/05/2020	€ 40.00	Buy
Kepler Cheuvreux	France	18/05/2020	€ 40.00	Buy
LifeSci Capital Alpha Series	US	14/05/2020	€ 41.00	Buy
Portzamparc	France	25/05/2020	€ 24.70	Buy
General recommendation: Buy		Average target price: € 35.50		

For full access to the reports, please directly contact the respective analysts listed on [Abivax's website](#).

Highly experienced Executive Committee



Prof. Hartmut Ehrlich, M.D.
Chief Executive Officer
Former Head of Global R&D,
Baxter BioScience

Baxter **SANDOZ** **Lilly**



Didier Blondel
Chief Financial Officer &
Board Secretary

SANOFI **sanofi pasteur MSD**
vaccines for life



Pierre Courteille
Pharmacist, MBA
Chief Commercial
Officer & VP, BD

sanofi pasteur **Guerbet**
Contrast for Life



Jérôme Denis
Ph.D.
VP, Process Dev. &
Manufacturing

IMA **LYONBIOPOLE**



Alexandra Pearce
Ph.D.
VP, Regulatory Affairs,
Quality, PV

AMGEN **Pfizer**



Paul Gineste
Pharm.D.
VP, Clinical
Operations

Boehringer Ingelheim **ALTANA**



Didier Scherrer
Ph.D.
VP, R&D

AstraZeneca



Jean-Marc Steens
M.D.
Chief Medical
Officer

ViiV Healthcare **gsk** **GlaxoSmithKline**



Prof. Jamal Tazi
Ph.D.
VP, Research

CJRS **W**

→ Competencies from discovery to global commercialization